

AD-A043 672

FLORIDA UNIV GAINESVILLE CENTER FOR MATHEMATICAL SYS--ETC F/G 12/1  
ON THE OBSERVABILITY OF POLYNOMIAL SYSTEMS.(U)  
1977 E D SONTAG


UNCLASSIFIED

AFOSR-TR-77-0917 AF-AFOSR-2268-72  
NL

| OF |  
AD  
A043672



END  
DATE  
FILMED  
9-77  
DDC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS E (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR-77-0917</b>	2. GOVT ACCESSION NO.	3. REPORT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>ON THE OBSERVABILITY OF POLYNOMIAL SYSTEMS</b>	5. TYPE OF REPORT & PERIOD COVERED <b>Interim / rept.</b>	
6. AUTHOR(s) <b>Eduardo D. Sontag</b>	7. PERFORMING ORG. REPORT NUMBER	
8. PERFORMING ORGANIZATION NAME AND ADDRESS <b>Center for Mathematical System Theory University of Florida Gainesville, FL 32611</b>	9. CONTRACT OR GRANT NUMBER(s) <b>AFOSR 72-2268</b>	
10. CONTROLLING OFFICE NAME AND ADDRESS <b>Air Force Office of Scientific Research/NM Bolling AFB DC 20332</b>	11. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>61102F 23041A1</b>	
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. REPORT DATE <b>1977</b>	
	14. NUMBER OF PAGES <b>8 pages.</b>	
	15. SECURITY CLASS. (of this report) <b>UNCLASSIFIED</b>	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) <b>✓ AF-AFOSR-2268-7231</b> <b>✓ DA-ARO-D-31-124-72-G114</b> Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES <b>Submitted to SIAM J. Control and Optimization.</b>		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <b>Nonlinear systems observability nonlinear filtering</b>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <b>Four natural notions of observability are compared in the case of systems described by polynomial difference equations. The main result states that, for a system having the standard property of (multiple-experiment initial-state) observability, the response to almost any (long-enough) input sequence is sufficient for final-state determination.</b>		

A043672

FILE COPY

DDC  
RECEIVED  
SEP 8 1977  
A

**AFOSR-TR- 77 - 0917**

ON THE OBSERVABILITY OF POLYNOMIAL SYSTEMS

by

Eduardo D. Sontag<sup>\*</sup>

Address until August 1977:


Center for Mathematical System Theory  
Department of Mathematics, Univ. of Florida  
Gainesville, FL 32611

after September 1977:

Department of Mathematics, Rutgers University  
New Brunswick, N.J. 08903

Approved for public release;  
distribution unlimited.

1977



-----  
\* This research was supported in part by US Army Research Grant DA-ARO-D-31-124-72-G114 and by US Air Force Grant 72-2268 through the Center for Mathematical System Theory, University of Florida, Gainesville, FL 32611, USA.

7180-77-8T-8E07A

**AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)**  
**NOTICE OF TRANSMITTAL TO DDC**  
This technical report has been reviewed and is  
approved for public release IAW AFR 190-12 (7b).  
Distribution is unlimited.  
**A. D. BLOSE**  
Technical Information Officer



# ABSTRACT

Four natural notions of observability are compared in the case of systems described by polynomial difference equations. The main result states that, for a system having the standard property of (multiple-experiment initial-state) observability, the response to almost any (long-enough) input sequence is sufficient for final-state determination.

APPROVED BY	
RTM	White Section <input checked="" type="checkbox"/>
RDC	Ext. Section <input type="checkbox"/>
WARRANTY	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION AVAILABILITY CODES	
ONE	AVAIL. AND. IN SPECIAL
A	

## 1. DEFINITIONS AND PROBLEM STATEMENT

We deal in this paper with discrete-time finite-dimensional nonlinear systems whose next-state and output maps are polynomial functions of previous states and inputs:

$$\Sigma \quad \begin{cases} x(t+1) = P(x(t), u(t)) \\ y(t) = h(x(t)), \end{cases} \quad t = 0, 1, 2, \dots,$$

where  $u(t)$ ,  $x(t)$ ,  $y(t)$  belong to algebraic sets  $U$ ,  $X$ ,  $Y$  over an arbitrary fixed infinite field  $k$ , and where  $P: X \times U \rightarrow X$  and  $h: X \rightarrow Y$  are polynomial maps. (An algebraic set is a set, in some affine space  $k^n$ , defined by polynomial equations  $\{Q_i(x_1, \dots, x_n) = 0\}$ .) We assume that  $U$  is irreducible, i.e.  $U$  cannot be expressed as a union of two proper algebraic subsets. (This last restriction is made purely for technical convenience; note that, in particular, any affine space  $U = k^m$  is irreducible.)

Such  $\Sigma$  were called polynomial systems in SONTAG and ROUCHALEAU[1975], where various properties were studied using elementary algebraic-geometric tools; all the algebraic-geometric concepts needed for this paper are summarized in that reference.

Denoting the extension of  $P$  to input sequences also by  $P: X \times U^* \rightarrow X$ , [for the empty sequence  $e$ ,  $P(x, e) = x$ ,] we can give the following

(1.1) DEFINITIONS. A polynomial system  $\Sigma$  is:

(a) observable iff for each pair of distinct states  $x, z$  in  $X$  there exists an input sequence  $w$  in  $U^*$  such that

$$h(P(x, w)) \neq h(P(z, w)).$$

(One says that  $w$  distinguishes between  $x$  and  $z$ .)

(b) single-experiment observable iff there is some  $r \geq 0$  and some (fixed) input sequence  $w = (u_1, \dots, u_r)$  such that for any  $x \neq z$  in  $X$ ,

$$h(P(x, u_1, \dots, u_t)) \neq h(P(z, u_1, \dots, u_t))$$

for some  $0 \leq t \leq r$ .

(c) final-state determinable iff there is some  $r \geq 0$  and some (fixed) input sequence  $w = (u_1, \dots, u_r)$  such that

(\*) for every pair of states  $x, z$  in  $X$ , either  
 $h(P(x, u_1, \dots, u_t)) \neq h(P(z, u_1, \dots, u_t))$  for some  
 $0 \leq t \leq r$  or  $P(x, w) = P(z, w)$ .

(d) final-state determinable by generic inputs iff there is some  $r \geq 0$   
 and a proper algebraic set  $F \subset U^r$  such that (\*) holds for each  $w$  not in  $F$ .

We wish to prove that the following are precisely the implications that hold:

$$(1.2) \quad (b) \Rightarrow (a) \Rightarrow (d) \Rightarrow (c).$$

We show below, via counterexamples (1.3), (1.4) and (1.5) respectively,  
 that (a)  $\nRightarrow$  (b), (d)  $\nRightarrow$  (a) and (c)  $\nRightarrow$  (d), while Theorem (2.5) shows that (a)  $\Rightarrow$  (d).

(1.3) EXAMPLE.  $\Sigma_1$  has  $X := k^2$ ,  $U=Y:=k$  and equations

$$\begin{cases} x_1(t+1) = 0, \\ x_2(t+1) = x_1(t)(u(t) - x_1(t)) \\ y(t) = x_2(t). \end{cases}$$

is not difficult to verify that  $\Sigma_1$  is observable, in fact even algebraically  
servable in the sense of SONTAG and ROUCHALEAU[1975]. But no single sequence  
 $w$  serves to distinguish every pair of initial states: let  $w = (u, w')$ , with  
 $u$  in  $U$ ; if  $u = 0$  then  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} -1 \\ 0 \end{pmatrix}$  are not distinguished by  $w$ , while  
 if  $u \neq 0$  then  $\begin{pmatrix} u \\ 0 \end{pmatrix}$  is indistinguishable from  $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$ .

(1.4) EXAMPLE.  $\Sigma_2$  has  $X = Y := k$ ,  $U$  arbitrary, and equations

$$\begin{cases} x(t+1) = 0 \\ y(t) = 0. \end{cases}$$

Clearly  $\Sigma_2$  is not observable, although (d) holds trivially with  $F = \text{empty}$ ,  $r = 1$ .

(1.5) EXAMPLE.  $\Sigma_3$  has  $X = U = Y := k$  and equations

$$\begin{cases} x(t+1) = x(t)u(t) \\ y(t) = 0. \end{cases}$$

The sequence (of length one)  $w = 0$  solves the final-state determination  
 problem. Comparing any nonzero state  $x$  with the zero state shows that  
 any sequence solving the final-state determination problem must have at  
 least one nonzero component, so no proper set  $F$  as in (d) can exist.



(1.6) DISCUSSION. (i) Note that (a) is the definition of observability standard in system theory, and that, trivially, (b) implies (a) and (d) implies (c). Property (c) assures the existence of an input sequence  $w$  such that, independently of the initial state  $x(0)$ , the final state  $x(r)$  can be determined from the output data  $y(0), \dots, y(r)$ . On the other hand, (d) guarantees that such a determination is theoretically possible with essentially any long-enough input sequence, i.e. in real-time system operation. [Of course, this determination is only insured in principle, and effective algorithms inverting the map  $(y(0), \dots, y(r)) \mapsto x(r)$  must still be developed prior to practical applications to nonlinear filtering problems.]

(ii) For linear systems, (a) is equivalent to (b) and (c) is equivalent to (d); see KALMAN, FALB and ARBIB[1969, Chapters 2 and 10].

(iii) The notions corresponding to (a), (b) and (c) have been studied in great detail for finite automata, beginning with MOORE[1956]; see GILL[1962] or CONWAY[1971]. In the automata literature one calls (a) a "diagnosing" and (c) a "homomorphism" problem, and one has precisely the implications  $(b) \Rightarrow (a) \Rightarrow (c)$ . [The automata case takes care of polynomial systems over a finite field  $k$ ; this allows us to restrict attention to infinite  $k$ , with the notational advantage of allowing identification of polynomials and polynomial functions.] Finiteness of the state set is crucial in the arguments involved in the automata results, but there appears to be no way of generalizing such arguments through a replacement of finiteness by finite-dimensionality: it can be shown, for instance, that in the context of systems defined by analytic (rather than polynomial) equations, (a) does no longer imply (c), say over  $k = \text{reals}$ . Thus, one needs to develop completely new arguments here.

(iv) For continuous-time systems, time-reversibility of differential equations implies that (b) is equivalent to (c); this helps perhaps to understand intuitively why (a), (b), (c) may be equivalent in that context, as shown recently by GRASSELLI and ISIDORI[1977] for the case of internally-bilinear continuous-time systems.

## 2. PROOF OF THE MAIN RESULT

(2.1) LEMMA. For any polynomial system  $F$  there exists an integer  $r \geq 0$  and a proper algebraic set  $F \subset U^r$  such that, for every  $w = (u_1, \dots, u_r)$  not in  $F$ , and for any  $x, z$  in  $X$ ,

$$h(P(x, u_1, \dots, u_t)) = h(P(z, u_1, \dots, u_t)), \quad t = 0, \dots, r$$

implies

$$P(x, w) \text{ is indistinguishable from } P(z, w).$$

PROOF. Since  $Y \subseteq k^p$  for some integer  $p$  and since a union of proper algebraic subsets of  $U^r$  is again a proper algebraic subset, it is sufficient to prove the lemma with  $Y = k$ .

Let  $s \geq 0$  be such that any pair of distinguishable states is already distinguished by inputs of length  $\leq s$  (SONTAG and ROUCHALEAU [1975, Corollary 7.3]).

For any algebraic set  $Z$ , let  $A(Z)$  denote the algebra of polynomial functions on  $Z$ . Let  $D$  be the direct limit of the sequence of  $k$ -algebras

$$A(U) \rightarrow \dots \rightarrow A(U^t) \rightarrow A(U^{t+1}) \rightarrow \dots,$$

where

$$A(U) \rightarrow A(U^{t+1}) = A(U^t) \otimes A(U): f \mapsto f \otimes 1.$$

Let  $K$  be the quotient field of  $D$  (which is an integral domain because  $U$  is irreducible).

Since  $Y = k$ , a polynomial map  $X \times X \times U^t \rightarrow Y$  is an element of  $A(X \times X) \otimes A(U^t)$ ; in particular the functions  $h_t$  defined by

$$h_t(x, z, u_1, \dots, u_t) := h(P(x, u_1, \dots, u_t)) - h(P(z, u_1, \dots, u_t))$$

are in  $A(X \times X) \otimes K$ . The latter is a finitely generated algebra over the field  $K$ , hence Noetherian. Thus there is some integer  $r$  such that all  $h_t$  are in the ideal of  $A(X \times X) \otimes K$  generated by  $h_0, \dots, h_r$ . In particular, there are therefore equations

$$(2.2) \quad ch_{r+j} = \sum_{t=0}^r a_{jt} h_t, \quad j = 1, \dots, s,$$

with all  $a_{jt}$  in  $A(X \times X) \otimes D$  and  $c$  a nonzero element of  $D$ . Since  $D$  is the union of the  $A(U^t)$ , there is some integer  $q$  such that all  $a_{jt}$  are in  $A(X \times X) \otimes A(U^q)$  and  $c$  is in  $A(U^q)$ . Without loss of generality, we shall assume that  $q \geq r + s$ .

Define the proper algebraic set

$$F := \left\{ (u_1, \dots, u_r) \text{ in } U^r \text{ such that } c(u_1, \dots, u_r, \dots, u_q) = 0 \text{ for all } (u_{r+1}, \dots, u_q) \right\}.$$

Claim:  $F$  satisfies the requirements of the lemma. Indeed, assume that  $\underline{w} = (u_1, \dots, u_r)$  is not in  $F$ . Take  $x, z$  in  $X$  such that  $h(P(x, u_1, \dots, u_t)) = h(P(z, u_1, \dots, u_t))$  for all  $t = 0, \dots, r$ , i.e.,

$$(2.3) \quad h_t(x, z, u_1, \dots, u_t) = 0, \quad t = 0, \dots, r.$$

Denote  $\underline{x} := P(x, \underline{w})$ ,  $\underline{z} := P(z, \underline{w})$ . It must be proved that  $\underline{x}, \underline{z}$  are indistinguishable.

Assume that  $\underline{x}, \underline{z}$  are distinguished by an input sequence  $v$ , which can be taken of length  $j$ ,  $0 \leq j \leq s$ , by definition of  $s$ . Let

$$F_1 := \left\{ w \text{ in } U^j \text{ such that } h_{r+j}(x, z, \underline{w}, w) = 0 \right\};$$

this is an algebraic set, proper because  $v$  is not in  $F_1$ . Let

$$F_2 := \left\{ w \text{ in } U^j \text{ such that } c(\underline{w}, w, w') = 0 \text{ for all } w' \text{ in } U^{q-r-j} \right\};$$

this is also an algebraic set, and it is proper because  $\underline{w}$  was taken not in  $F_2$ .

It follows that  $F_1 \cup F_2$  is also a proper algebraic set. Let then  $w$  be in neither  $F_1$  nor  $F_2$ . Then  $c(\underline{w}, w, w') \neq 0$  for some  $w'$ , so

$$(2.4) \quad c(\underline{w}, w, w') h_{r+j}(x, z, \underline{w}, w) \neq 0.$$

But (2.2), (2.3) and (2.4) taken together are contradictory. □

(2.5) THEOREM. Observability implies, for polynomial systems, final-state determinability with generic inputs.

PROOF. Immediate from the lemma. □

(2.6) REMARK. As shown in SONTAG[1976], canonical realizations  $\Sigma_f$  of polynomial response maps are not, in general, polynomial systems. So the Theorem above is not applicable directly ( $A(X_f)$  is not Noetherian). However, if  $f$  admits a polynomial



realization  $\Sigma$ , then the reachable states of  $\Sigma_f$  form a set which is a quotient of the reachable states of  $\Sigma$ . Then lemma (2.1) can be applied to  $\Sigma$ , implying that the reachable part of  $\Sigma_f$  does satisfy (2.2).

## REFERENCES

J. H. CONWAY

[1971] Regular Algebra and Finite Machines, Chapman and Hall, London.

A. GILL

[1962] Introduction to the Theory of Finite-State Machines, McGraw-Hill, New York.

O. M. GRASSELLI and A. ISIDORI

[1977] "Deterministic state reconstruction and reachability of bilinear control processes", Proc. JACC, San Francisco, June 22-25, 1977.

R. E. KALMAN, P. L. FALB and M. A. ARBIB

[1969] Topics in Mathematical System Theory, McGraw-Hill, New York.

E. F. MOORE

[1956] "Gedanken experiments on sequential machines", in Automata Studies, Princeton University Press, Princeton, N.J..

E. D. SONTAG

[1976] "On the Internal Realization of Polynomial Response Maps", Doctoral Dissertation, University of Florida.

E. D. SONTAG and Y. ROUCHALEAU

[1975] "On discrete-time polynomial systems", CNR-CISM Symposium on Algebraic System Theory, Udine, Italy. Appeared in revised form in J.Nonlinear Analysis, Methods, Theory and Applications, 1(1976):55-64.



